
Homework 11

- Let M be the subspace of \mathbb{R}^4 spanned by $v_1 = (1, 1, 0, 0)$; $v_2 = (0, 1, 1, 0)$; $v_3 = (0, 0, 1, 1)$; and $v_4 = (1, 0, 0, 1)$.
 - Use the Gram-Schmidt orthogonalization process to find an orthonormal basis for M .
 - Extend the basis you found in part (a) to an orthonormal basis for all of \mathbb{R}^4 .
 - Is $w = (1, -1, 0, 1)$ in M ? (Justify your answer.)
- Let $C = \begin{pmatrix} 1 & 0 & 3 & -1 \\ 0 & 1 & 1 & 2 \\ 2 & -1 & 5 & -4 \end{pmatrix}$ (Compare with problems from Homework 7).
 - Find an orthonormal basis for the nullspace of C .
 - Find an orthonormal basis for the range of C .
 - Find the angles between the vectors you found in parts (a) and (b).
 - Find an orthonormal basis for the nullspace of C' .
 - Find an orthonormal basis for the range of C .
 - What do you notice about your answers to parts (d) and (e).
- Let M be the subspace of \mathbb{R}^4 spanned by $(2, 1, 0, -1)$, $(1, 1, 1, 0)$, and $(-1, 0, 1, 1)$. Find w in M and u in M^\perp so that $w + u = (3, 1, 0, 0)$.
- Let \mathcal{W} be the subspace of \mathbb{R}^5 spanned by $(1, 1, 0, -1, 1)$, $(0, 2, -1, 1, 1)$, and $(-1, 1, 1, -2, 1)$. Find w in \mathcal{W} and u in \mathcal{W}^\perp so that $w + u = (1, 1, -1, 1, 1)$.
- The matrix B is an 8×11 matrix and the dimension of $\mathcal{N}(B)$, the nullspace of B , is 5.
 - What is the dimension of $\mathcal{R}(B)$, the range of B ?
 - What is the dimension of $\mathcal{R}(B')$, the range of B' ?
 - What is the dimension of $\mathcal{N}(B')$, the nullspace of B' ?
 - What is the dimension of $\mathcal{R}(B')^\perp$, the orthogonal complement of $\mathcal{R}(B')$?
- An $n \times n$ matrix U is called *unitary* if $U' = U^{-1}$.
 - Show U is unitary if and only if its columns form an orthonormal basis for \mathbb{C}^n .
 - Show that if U is unitary, then U^{-1} is also unitary.
 - Show that if U is a real unitary matrix, the map $x \mapsto Ux$ is a rigid motion of \mathbb{R}^n by showing that if U is unitary and v and w are vectors in \mathbb{R}^n , then
 - $\langle Uv, Uw \rangle = \langle v, w \rangle$,
 - $\|Uv\| = \|v\|$,
 - the angle between Uv and Uw is the same as the angle between v and w .
- Find the general solution of the system *and find three explicit solutions*

$$\begin{cases} a - b + 2c & + e = 0 \\ -a + 2b - 3c + 2d - e & = 1 \\ a - 2b + 4c - d + 2e & = -1 \\ -2a + 3b - 2c + 5d + e & = 1 \end{cases}$$
- Let M be the subspace of \mathbb{R}^5 spanned by $w_1 = (1, 1, 0, -1, 1)$, $w_2 = (1, 0, 2, 0, -1)$, and $w_3 = (0, 1, 1, 1, -1)$. Find vectors w in M and u in M^\perp so that $(1, 0, 0, 1, 1) = w + u$.
- Let M be the subspace of \mathbb{R}^5 spanned by $(2, 1, 0, -1, 1)$, $(1, 1, 1, 0, 2)$, and $(-1, 0, 1, 1, 1)$. Find w in M and u in M^\perp so that $w + u = (3, 1, 0, 0, -1)$.
