## February 8

- \* 26. (a) Let  $v_1 = (1, -1)$ ,  $v_2 = (2, -1)$ , and  $v_3 = (-3, 2)$ . Let  $w_1 = (1, 0)$ ,  $w_2 = (0, 1)$ , and  $w_3 = (1, 1)$ . Is there a linear transformation T from  $\mathbb{R}^2$  into  $\mathbb{R}^2$  so that  $T(v_j) = w_j$  for j = 1, 2, 3?
  - (b) Give necessary and sufficient conditions on  $u_1$ ,  $u_2$ , and  $u_3$  in  $\mathbb{R}^2$  so that there is a linear transformation T from  $\mathbb{R}^2$  into  $\mathbb{R}^2$  for which  $T(u_j) = w_j$  for j = 1, 2, 3?
- \* 27. Let V be an n-dimensional vector space over F and let T be a linear transformation for which the range of T and the null space of T are the same subspace.
  - (a) Prove that n is even.
  - (b) Give an example of a linear transformation S acting on  $\mathbb{R}^4$  for which the range of S and the null space of S are spanned by  $v_1 = (1, 0, 1, -1)$ ,  $v_1 = (1, 1, -1, 2)$ , and  $v_3 = (3, 1, 1, 0)$ .
- \* 28. Let V be a finite dimensional vector space over the field F and let T be a linear transformation for which the rank of  $T^2$  and the rank of T are the same. Prove that the intersection of the range of T and the null space of T is the zero subspace (0).
- \*\* **29.** If  $p(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_k x^k$  where the  $a_j$  are real numbers, and A is an  $n \times n$  real matrix, we define p(A) to be  $a_0I + a_1A + a_2A^2 + \dots + a_kA^k$ .
  - (a) What is the dimension of the vector space of  $3 \times 3$  real matrices?
  - (b) Prove that for any  $3 \times 3$  real matrix A, there is a polynomial p so that p(A) = 0.
- \* 30. For n a positive integer, let T be a linear transformation of the vector space  $F^n$  into the space  $F^m$ . Let A be the matrix for T with respect to the standard bases for  $F^n$  and  $F^m$ . Let W be the subspace of  $F^m$  spanned by the columns of A. State one or two relationships between W and T.
- \* **31.** Let *T* be the linear operator on  $\mathbb{R}^3$  defined by  $T(x_1, x_2, x_3) = (3x_1 + x_3, -2x_1 + x_2, -x_1 + 2x_2 + 4x_3)$ (1)
  - (a) What is the matrix for T in the standard basis?
  - (b) Find the matrix for T relative to the basis

 $v_1 = (1, 0, 1)$   $v_2 = (-1, 2, 1)$   $v_3 = (2, 1, 1)$ 

(c) Prove that T is invertible and find an expression for the transformation  $T^{-1}$  like the one in Equation (1).

\*\* **32.** Let V be an n-dimensional vector space over the field F and let  $\mathcal{B} = \{v_1, v_2, \dots, v_n\}$  be an ordered basis for V.

- (a) Let T be the (unique) linear transformation that satisfies  $Tv_j = v_{j+1}$  for  $j = 1, 2, \dots, n-1$  and  $Tv_n = 0$ . Find the matrix for T with respect to basis  $\mathcal{B}$  above.
- (b) Prove that  $T^n = 0$  but  $T^{n-1}$  is not the zero transformation.
- (c) Let S be an operator on V for which  $S^n = 0$  but  $S^{n-1} \neq 0$ . Prove that there is an ordered basis  $\mathcal{B}'$  for V such that the matrix for S with respect to the basis  $\mathcal{B}'$  is the same as the matrix from part (a) of this exercise.