



Presentation Abstract

Program#/Poster#: 298.17/JJJ20

Title: Sympatho-respiratory coupling: Brainstem mechanisms

Location: Halls B-H

Presentation Time: Sunday, Nov 14, 2010, 1:00 PM - 2:00 PM

Authors: \***Y. I. MOLKOV**<sup>1</sup>, D. B. ZOCCAL<sup>2</sup>, D. J. A. MORAES<sup>2</sup>, D. M. BAEKEY<sup>3</sup>, T. E. DICK<sup>4</sup>, B. H. MACHADO<sup>2</sup>, J. F. R. PATON<sup>5</sup>, I. A. RYBAK<sup>1</sup>;

<sup>1</sup>Dept. of Neurobio. and Anat., Drexel Univ. Col. of Med., Philadelphia, PA; <sup>2</sup>Dept. Physiol., Univ. of São Paulo, Ribeirão Preto, São Paulo, Brazil; <sup>3</sup>Dept. Med., <sup>4</sup>Dept. Med. & Neurosci, Case Western Reserve Univ., Cleveland, OH; <sup>5</sup>Dept. Physiol. & Pharmacol., Univ. of Bristol, Bristol Heart Inst., Bristol, United Kingdom

Abstract: Hypertension, like that resulting from sleep-disordered breathing, is associated with elevated sympathetic nerve activity (SNA), which was found to be respiratory phase-dependent. Thus, hypertension may depend on specific interactions between the respiratory and sympathetic neural networks. Conditioning animals with chronic intermittent hypoxia (CIH) increases the late-expiratory (late-E) component in the SNA. In contrast, in naïve animals during eupnea SNA is typically most active in inspiration/post-inspiration and least active during late-E. Further, our recent studies *in situ* demonstrate that ponto-medullary transection significantly attenuates SNA respiratory modulation. Also, baro-stimulation facilitates post-inspiratory (post-I) activity in the Bötzing Complex. However, the sources of respiratory modulation of SNA and the mechanisms responsible for the coupling between respiratory and sympathetic activities remain undefined. To elucidate the mechanisms underlying these phenomena we extended our computational model of the brainstem respiratory network by incorporating neural populations representing rostral and caudal ventrolateral medulla (RVLM and CVLM). In this model, the RVLM and CVLM populations receive inputs from neural populations of the pons, Bötzing and pre-Bötzing Complexes and retrotrapezoid nucleus/parafacial respiratory group (RTN/pFRG), which together provide the respiratory modulation of SNA. Our simulations suggest that CIH-conditioning may evoke a late-E modulation of SNA similar to that produced by hypercapnia. This suggestion was experimentally confirmed *in situ*. We found that CIH-conditioned rats under normal conditions exhibit synchronized late-E discharges in the abdominal nerve (AbN) and SNA, which were similar to the late-E activity observed in both nerves in control rats during hypercapnia. With progressive increase of CO<sub>2</sub> level in control rats, late-E discharges (in both AbN and SNA) appeared at a lower rate relative to phrenic nerve (PN), i.e. many respiratory cycles were skipped, and then quantally accelerated until they occurred in every cycle. The SNA late-E discharges appeared during the same breathing cycles as AbN late-E bursts. Similar behavior was observed in CIH-conditioned rats at normal CO<sub>2</sub>. We hypothesize that the mechanism for emergence of late-E modulation of SNA in CIH-conditioned animals is associated with an increase in the CO<sub>2</sub> sensitivity of RTN/pFRG neurons as a consequence of intermittent hypocapnia resulting from tachypneic response to hypoxia. This suggestion is supported by a lower apnea threshold found in CIH-conditioned rats relative to naïve ones.

Disclosures: **Y.I. Molkov**, None; **D.B. Zoccal**, None; **D.J.A. Moraes**, None; **D.M. Baekey**, None; **T.E. Dick**, None; **B.H. Machado**, None; **J.F.R. Paton**, None; **I.A. Rybak**, None.

Keyword(s): RESPIRATION  
HYPERTENSION  
SYMPATHETIC

Support: NIH Grant R33 HL087377  
NIH Grant R33 HL087379  
NIH Grant R01 HL090554

NIH Grant R01 HL090554

NIH Grant R01 NS057815

AHA SDG 0735037N

[Authors]. [Abstract Title]. Program No. XXX.XX. 2010 Neuroscience Meeting Planner. San Diego, CA: Society for Neuroscience, 2010. Online.

2010 Copyright by the Society for Neuroscience all rights reserved. Permission to republish any abstract or part of any abstract in any form must be obtained in writing by SfN office prior to publication.