

**Exercises to Accompany**  
**Lectures on Composition Operators on**  
**Spaces of Analytic Functions**

Carl C. Cowen

IUPUI

(Indiana University Purdue University Indianapolis)

GSMAA Workshop on Operator Theory, Helsinki, 7–11 May 2012

**From Monday, May 7**

**Exercise 1:** A proof of the assertion that  $L^1([0, 1])$  is *not* a functional Banach space with  $X = [0, 1]$  and the obvious identification of integrable functions as vectors in  $L^1$ :

The continuous functions on  $[0, 1]$  are a dense subset of  $L^1([0, 1])$  in the  $L^1$  norm and if  $f$  is a continuous function on  $[0, 1]$  then  $f(1/2)$  is defined. Even more, for a continuous function on  $[0, 1]$ , the value of  $f$  at  $x = 1/2$  cannot be changed and still have  $f$  continuous at  $x = 1/2$ .

Show that, considering the continuous functions on  $[0, 1]$  as a subset of  $L^1([0, 1])$ , the linear functional on this subset  $f \mapsto f(1/2)$  is *not* bounded.

**Exercise 2:** Prove that the Bergman space is a Hilbert space, that is, that it is a *complete* inner product space. Equivalently, since it is obvious that  $A^2(\mathbb{D}) \subset L^2(\mathbb{D})$  and we know  $L^2$ , it is enough to show that  $A^2(\mathbb{D})$  is a closed subset of  $L^2(\mathbb{D})$ . That is, show that if  $f_n$  is a sequence of functions in  $A^2(\mathbb{D})$ , and  $\lim_{n \rightarrow \infty} f_n = f$  in  $L^2$ , then actually  $f$  is analytic also and is  $A^2(\mathbb{D})$ .

**Exercise 3:** Just as for  $H^2(\mathbb{D})$ , we want another way to think about  $A^2(\mathbb{D})$ .

- (a) Show that the set  $\{z^n\}_{n=0}^{\infty}$  is an *orthogonal* basis for  $A^2(\mathbb{D})$ .
- (b) Find the norm of  $z^n$  in  $A^2(\mathbb{D})$  for each non-negative integer  $n$ .
- (c) Find a condition (\*) on the coefficients  $a_n$  so that if  $f$  is an analytic function on the disk with  $f(z) = \sum_{n=0}^{\infty} a_n z^n$ , then  $f$  is in  $A^2(\mathbb{D})$  if and only if (\*).
- (d) Use the ideas of (a)–(c) to show that for  $\alpha$  in the disk, the function  $K_\alpha$  in  $A^2(\mathbb{D})$  so that  $\langle f, K_\alpha \rangle = f(\alpha)$  for every  $f$  in  $A^2(\mathbb{D})$  is

$$K_\alpha(z) = \frac{1}{(1 - \bar{\alpha}z)^2}$$

**Exercise 4:** Find all the fixed points of the listed functions and their derivatives there. Then find the Denjoy–Wolff point. ( $\sqrt{\phantom{x}}$  means the branch of the square root that is positive on the positive axis.)

(a)  $\varphi(z) = \exp((z+1)/(z-1))$

(b)  $\varphi(z) = \left(\frac{z+1/3}{1+z/3}\right)^2$

(c)  $\varphi(z) = w^{-1}(\psi(w(z)))$  where  $w(z) = (1+z)/(1-z)$  maps the disk to the right halfplane and  $\psi(w) = \sqrt{4w^2+3}$  maps the right half plane to itself.

(d)  $\varphi(z) = \frac{1+z+2\sqrt{1-z^2}}{3-z+2\sqrt{1-z^2}}$

### From Wednesday, May 9

**Exercise 5:** For the maps in Exercise 4, find the Case of  $\varphi$ , that is, decide, for each  $\varphi$ , if  $\varphi$  is in the plane dilation, half-plane dilation, plane translation, or half-plane translation case.

**Exercise 6:** For the maps in Exercise 4, find the spectrum of  $C_\varphi$  as an operator on  $H^2$  if you can, or to the extent that you can.

### From Thursday, May 10

**Exercise 7:** Consider the map  $\varphi(z) = z/2 + z^2/3$ .

(a) Show that  $\varphi$  maps the disk into the disk.

(b) Explain why  $C_\varphi$  is compact on  $H^2$ .

(c) Find the spectrum of  $C_\varphi$ .

(d) According to the theory, the eigenvectors are multiples of powers of the Koenigs' function  $\sigma$ , which is also the map in the model for the function  $\varphi$ . In this case, we know  $\sigma(\varphi(z)) = \lambda\sigma(z)$  and  $0 < |\lambda| < 1$ , and this  $|\lambda|$  is the largest possible eigenvalue, with  $|\lambda| < 1$ . Find  $\lambda$  in this case, and find the first 7 Taylor coefficients, i.e. up to  $a_6$ , the coefficient of  $z^6$ , of  $\sigma$  explicitly. Looking at these Taylor coefficients, can you find  $\sigma$  explicitly, that is, can you guess the rest of the coefficients and write down  $\sigma$  as an elementary function?

**Exercise 8:** In the Theorem for the model for iteration in the case in which the Denjoy-Wolff point,  $a$ , is inside the disk, the hypothesis is  $\varphi'(a) \neq 0$ . Using  $\varphi(z) = z^2$ , which has Denjoy-Wolff point  $a = 0$  and  $\varphi'(a) = \varphi'(0) = 0$ , explain why the hypothesis is what it is by finding possible analytic functions  $f$  and numbers  $\lambda$  so that  $f(\varphi(z)) = \lambda f(z)$  in a neighborhood of 0.

**Exercise 9:** If  $\varphi$  maps the disk into itself and has Denjoy-Wolff point 1 with  $\varphi'(1) = .5$ , the theory says that the inductively defined sequence  $z_{n+1} = \varphi(z_n)$  starting with any point  $z_0$  in the disk is an interpolating sequence. For the function  $\varphi(z) = .5z + .5$ , find  $z_n$  explicitly satisfying  $z_{n+1} = \varphi(z_n)$  starting with  $z_0 = 0$ . Show that, at least,  $\{z_n\}$  is a Blaschke sequence, that is, that  $\sum(1 - |z_n|) < \infty$ , so that there are analytic functions  $f$  with  $f(z_n) = 0$  for all  $n$  but  $f$  is not the zero function.

**Exercise 10:** Let  $\varphi$  be an analytic function mapping the unit disk into itself, with  $\varphi(1) = 1$  and  $\varphi'(1) = s$  where  $0 < s < 1$ . According to the theory,  $\varphi$  is in the half-plane dilation case and there is  $\sigma$  analytic, mapping  $\mathbb{D}$  into the right half plane  $H_+ = \{z : \operatorname{Re} z > 0\}$  where  $\Phi(w) = sw$  and  $\Phi \circ \sigma = \sigma \circ \varphi$ . Suppose, in addition,  $\varphi$  is real on the real axis. Using the functions  $\tilde{\varphi}(z) = \overline{\varphi(\bar{z})}$ ,  $\tilde{\Phi}(z) = \overline{\Phi(\bar{z})}$ , and  $\tilde{\sigma}(z) = \overline{\sigma(\bar{z})}$ , show that  $\sigma$  is real on the real axis as well.

**Exercise 11:** Let  $\varphi$  be an analytic function mapping the unit disk into itself. If  $\Phi$  is an automorphism mapping  $\Omega$  onto itself, and  $\sigma$  maps the disk into  $\Omega$  such that  $\Phi \circ \sigma = \sigma \circ \varphi$ , then it must be the case that  $\sigma$  maps the fixed points of  $\varphi$  to the fixed points of  $\Phi$  in such a way that the attracting fixed point of  $\varphi$ , the Denjoy-Wolff point of  $\varphi$ , is mapped to the attracting fixed point of  $\Phi$  and the other fixed points of  $\varphi$  are mapped to the other fixed point of  $\Phi$ , or at least the iterates of  $\varphi$  leaving a fixed point of  $\varphi$  are mapped to iterates of  $\Phi$  leaving a fixed point of  $\Phi$ . Moreover, if  $\varphi$  maps a point of the circle to the circle, then  $\sigma$  must map that point to a point of the boundary of  $\sigma(\mathbb{D})$  that  $\Phi$  maps to a point of the boundary of  $\sigma(\mathbb{D})$ . Use these ideas to draw a connected, simply connected domain  $U$  in the complex plane that  $U$  contains 0, so that  $z$  in  $U$  implies  $z/2$  is in  $U$ , and so that if  $\sigma$  is the Riemann map of  $\mathbb{D}$  onto  $U$  that takes 0 to 0 and has  $\sigma'(0) > 0$ , then using  $\Phi(w) = w/2$ , the map  $\varphi(z) = \sigma^{-1}(\Phi(\sigma(z)))$  has  $\varphi(0) = 0$ ,  $\varphi'(0) = 1/2$ , and 1 and  $-1$  as fixed points of  $\varphi$ . Can you describe explicitly, a map,  $\varphi$ , of  $\mathbb{D}$  into  $\mathbb{D}$  with  $\varphi(0) = 0$ ,  $\varphi'(0) = 1/2$ ,  $\varphi(1) = 1$ , and  $\varphi(-1) = -1$ ?